



SAGINAW COUNTY  
COMMUNITY MENTAL  
HEALTH AUTHORITY

# An Overview of Trauma and Trauma-Informed Care 2024

Mary Baukus, MSW, LMSW, CCTP

[Mary.Baukus@sccmha.org](mailto:Mary.Baukus@sccmha.org)

989-272-7372



# Objectives

- Identify important trauma and trauma informed care definitions
- Identify and describe different types of trauma
- Understand and identify trauma symptoms and responses for adults and children
- Describe Trauma informed care and understand its importance
- Understand Re-traumatization and secondary trauma and how to prevent it



SAGINAW COUNTY  
COMMUNITY MENTAL  
HEALTH AUTHORITY

# A Brief Introduction

## Trauma-Informed Care

Introduction



“A trauma survivor who seeks services may interact with a dozen individuals before actually sitting down with a clinician trained to provide trauma services. A woman will have to make an appointment and speak with a receptionist. A man will enter the agency and walk past a security guard or maintenance worker. A family may stop for a snack at a hospital cafeteria. Once they are in the agency, they may encounter office workers, intake personnel, trainees, and anonymous clinicians. Any of these individuals has the opportunity to make a consumer’s visit to the service agency inviting or terrifying.” (Harris & Fallo, 2001, p 7)



# Important Definitions

- **Trauma:** defined as a psychologically distressing event that is outside the range of usual human experience, often involving a sense of intense fear, terror, or helplessness that creates significant and lasting damage to a person's mental, physical and emotional growth.
- **Complex Trauma:** can occur when an individual has been exposed over time to persistent abuse, neglect, violence or abandonment, especially as a child. The person may have experienced multiple traumas, and the complex trauma is often worse when the perpetrator was someone close to the child.
- **Re-traumatization:** re-traumatization entails replication of the event(s) or dynamics of an original trauma which triggers a response associated with the original trauma.
- **Secondary Trauma:** A state experienced by those helping people in distress; it is an extreme state of tension and preoccupation with the suffering of those being helped to the degree that it is traumatizing for the helper. It can occur after one difficult incident or prolonged incidences of trauma exposure. Often symptoms mimic post traumatic stress disorder symptoms.
- **Compassion Fatigue:** also known as Vicarious trauma, is a permanent change in the provider resulting from empathic engagement with a consumer's traumatic background. Before Vicarious Trauma comes a secondary traumatic stress incident. The response is cumulative. It is much more pervasive than burnout even though the symptoms at times are similar.



# Adverse Childhood Experiences (ACE) Study

- Co-PIs: Robert F. Anda, MD (CDC) and Vincent J. Felitti, MD (Kaiser Permanente)
- HMO enrollees, average age 59, interviewed about difficult childhood experiences.
- Test consists of 10 questions regarding adverse childhood experiences.
- More than 17,000 interviewed.
- Staggering results, many since replicated



SAGINAW COUNTY  
COMMUNITY MENTAL  
HEALTH AUTHORITY

# ACES & TIC

**THE RISE OF  
TRAUMA-INFORMED CARE —  
FROM SAN DIEGO TO  
KANSAS CITY**

▶ ▶▶ 🔊 0:00 / 2:13





SAGINAW COUNTY  
COMMUNITY MENTAL  
HEALTH AUTHORITY

# The 10 ACEs the researchers measured

- Physical, sexual and verbal abuse
- Physical and emotional neglect
- A family member who is:
  - depressed or diagnosed with other mental illness
  - addicted to alcohol or another substance
  - in prison
- Witnessing a mother being abused
- Losing a parent to separation, divorce or other reason
- **See next slide.**
- **If you are comfortable doing so, please take a few minutes to fill out the ACE questionnaire for your own information.**



# ACE Questionnaire

## Adverse Childhood Experience (ACE) Questionnaire Finding your ACE Score ra hbr 10 24 06

While you were growing up, during your first 18 years of life:

1. Did a parent or other adult in the household **often** ...  
Swear at you, insult you, put you down, or humiliate you?  
**or**  
Act in a way that made you afraid that you might be physically hurt?  
Yes No If yes enter 1 \_\_\_\_\_
2. Did a parent or other adult in the household **often** ...  
Push, grab, slap, or throw something at you?  
**or**  
Ever hit you so hard that you had marks or were injured?  
Yes No If yes enter 1 \_\_\_\_\_
3. Did an adult or person at least 5 years older than you ever...  
Touch or fondle you or have you touch their body in a sexual way?  
**or**  
Try to or actually have oral, anal, or vaginal sex with you?  
Yes No If yes enter 1 \_\_\_\_\_
4. Did you **often** feel that ...  
No one in your family loved you or thought you were important or special?  
**or**  
Your family didn't look out for each other, feel close to each other, or support each other?  
Yes No If yes enter 1 \_\_\_\_\_
5. Did you **often** feel that ...  
You didn't have enough to eat, had to wear dirty clothes, and had no one to protect you?  
**or**  
Your parents were too drunk or high to take care of you or take you to the doctor if you needed it?  
Yes No If yes enter 1 \_\_\_\_\_
6. Were your parents **ever** separated or divorced?  
Yes No If yes enter 1 \_\_\_\_\_
7. Was your mother or stepmother:  
**Often** pushed, grabbed, slapped, or had something thrown at her?  
**or**  
**Sometimes or often** kicked, bitten, hit with a fist, or hit with something hard?  
**or**  
Ever repeatedly hit over at least a few minutes or threatened with a gun or knife?  
Yes No If yes enter 1 \_\_\_\_\_
8. Did you live with anyone who was a problem drinker or alcoholic or who used street drugs?  
Yes No If yes enter 1 \_\_\_\_\_
9. Was a household member depressed or mentally ill or did a household member attempt suicide?  
Yes No If yes enter 1 \_\_\_\_\_
10. Did a household member go to prison?  
Yes No If yes enter 1 \_\_\_\_\_

Now add up your "Yes" answers: \_\_\_\_\_ This is your ACE Score



# ACEs Study



- 1 in 16 smokes;  
1 in 14 has heart disease
- 1 in 69 misuses alcohol;  
**1 in 480 uses IV drugs**
- **1 in 96 has attempted suicide**


- **With 3 ACEs**, 1 in 9 smokes, 1 in 7 heart disease
- 1 in 9 misuses alcohol, 1 in 43 uses IV drugs
- 1 in 10 has attempted suicide

- **With 7+ ACEs**, 1 in 6 smokes, 1 in 6 has heart disease
- 1 in 6 misuses alcohol, **1 in 30 uses IV drugs**
- **1 in 5 has attempted suicide**



SAGINAW COUNTY  
COMMUNITY MENTAL  
HEALTH AUTHORITY

# Trauma and the Brain



Trauma and the Brain



# Types of Traumatic Events

- Sexual abuse
- Physical abuse
- Abandonment, betrayal of trust (Such as abuse by a caregiver)
- Neglect
- The death of a loved one
- Life threatening illness of a caregiver
- Witnessing domestic violence
- Bullying
- Military combat
- Serious automobile accidents
- Life-threatening health situations and/or painful medical procedures
- Witnessing police activity or having a close relative incarcerated
- Personal Incarceration
- Police Violence
- House fires or other reasons for loss of home or dwelling (Safe place)
- Life-threatening natural disasters
- Acts of threats or terrorism
- Living in poverty
- Being in special education (Children)-results in bullying



# Trauma Prevalence

- An estimated **70%** of adults in the United States have experienced a traumatic event at least once in their lives.
- Up to **90%** of public mental health clients in have been exposed to trauma (National Council for Mental Well-Being, 2022)
- On average an individual will experience at least **5** traumatic experiences in their lifetime.
- More than **90%** of individuals with developmental disabilities will experience some form of physical or sexual abuse in their lifetimes and **49%** will experience multiple instances of abuse.
- Traumatic events are associated with an increased likelihood of trauma survivors to develop dementia in older adulthood.





SAGINAW COUNTY  
COMMUNITY MENTAL  
HEALTH AUTHORITY

## Trauma in the LGBTQ Population

- LGBTQ people who experience microtraumas may not meet diagnosable criteria for PTSD yet suffer tremendously from minority stressors such as from internalized phobia, rejection sensitivity, marginalization, and discrimination both in their personal life and health care settings.
- 37% of LGBTQ youth report high levels of trauma symptoms and 60% report low to moderate levels of trauma symptoms.
- Symptoms of trauma are higher amongst BIPOC LGBTQ youth.





# Trauma and Substance Use Disorder



- 18% to 53.7% of patients who seek treatment for substance use disorder (SUD) present a history of physical abuse.
- 9.2% to 22.7% have experienced lifetime sexual abuse.
- Most experienced the traumatic event in childhood or adolescence.



SAGINAW COUNTY  
COMMUNITY MENTAL  
HEALTH AUTHORITY

# Veterans: PTSD & SUD

- More than 2 of 10 Veterans with Post Traumatic Stress Disorder also have SUD.
- Almost 1 out of every 3 Veterans seeking treatment for SUD also has PTSD.
- The number of Veterans who smoke (nicotine) is almost double for those with PTSD (about 6 of 10) versus those without a PTSD diagnosis (3 of 10).





# Common Trauma Responses-Adults

## ***Physical:***

- Sleep disturbances- nightmares, insomnia, over-sleeping
- Chronic health conditions
- Musculoskeletal pains
- Somatization
  - emotional disturbances manifesting themselves in the body
  - Often physical symptoms have no explainable cause-a full physical work up should still be done

## ***Emotional:***

- Depressive Symptoms
- Flashbacks
- Suicidal Ideation
- Self-harm-way of grounding flashbacks
- Withdrawal from others
- Intense fear/anxiety-Hyperarousal (Also called Hypervigilance)
- Exaggerated Startle Response
- Excessive worry

## ***Relational:***

- Boundary Problems
- Sexual problems-promiscuity or withdrawal from sex
- Loss of friendships-Withdrawal from others





# Common Trauma Responses in Children

- Irritability
- ADHD type symptoms
- Fear
- Anxiety
- Attachment difficulties
- Boundary problems
- Lack of Trust
- Hypervigilance
- Tempter Tantrums
- Insomnia/ nightmares
- Enuresis/ Encopresis
- Isolation
- Fear of strangers or being left alone





# What is Trauma-Informed Care (TIC)

- Trauma-Informed Care (TIC) is an approach to the delivery of behavioral health services that includes an understanding of trauma and an awareness of the impact that it can have across settings, services, and populations.
- The primary purpose of TIC services is not the treatment of trauma. The purpose is to approach the person who has needs due to their trauma history in a sensitive, caring, and welcoming way.
- The idea is to use **universal trauma precautions**. This approach always presumes the possibility that an individual one encounters, whether a consumer, visitor, or staff member, may have a trauma history.



SAGINAW COUNTY  
COMMUNITY MENTAL  
HEALTH AUTHORITY

# What is Trauma-Informed Care (TIC)





# The Core Principles of a Trauma-Informed System of Care

- **Safety:** Ensuring physical and emotional safety of staff members, visitors, and persons served by providing a safe, calm, and secure environment with supportive care to ensure the physical and emotional safety of everyone.
- **Trustworthiness:** Maximizing trustworthiness, making tasks clear, and maintaining appropriate boundaries.
- **Choice:** Supporting staff member and consumer voice, choice, and control as well as self-advocacy.
- **Collaboration:** Maximizing collaboration and sharing of power with consumers through the development of healing, hopeful, honest, and trustworthy relationships.



# The Core Principles of a Trauma-Informed System of Care


- **Empowerment:** Prioritizing staff member and consumer empowerment and skill building through recovery-oriented, choice-driven, trauma specific services and supports.
- **System-wide understanding** of the meaning of trauma, the prevalence, and the impact of trauma on individuals.
- **Cultural Humility and Responsiveness:** The ability to understand, communicate with, and effectively interact with people across cultures. Biases and stereotypes (e.g., based on race, ethnicity, sexual orientation, age, geography) and historical trauma are recognized and addressed.



SAGINAW COUNTY  
COMMUNITY MENTAL  
HEALTH AUTHORITY

# Why Trauma-Informed Care?

- A trauma-informed organization:
  - Increases safety for all
  - Improves social environment in a way that improves relationships for all
  - Cares for the caregivers
  - Improves quality of services
  - Reduces negative encounters and events
  - Creates a community of hope, wellness and recovery
  - Increases success and satisfaction at work
  - Promotes organizational wellness
  - Helps to ensure consumers come back for services





# Trauma-Informed Care Basic Assumptions

- Trauma shapes the survivor's basic beliefs about identity, world view and spirituality, or meaning-making.
- **Trauma is something that happened to the individual not something that is wrong with them.**
- The four most important things a worker has to offer a survivor is Respect, Information, Connection and Hope (RICH).
- Workers need support from one another, including (RICH).
- Working with survivors affects the person of the helper as well.





# Trauma-Informed Treatment Objectives and Strategies

- Establish safety
- Prevent Re-traumatization
- Provide Psychoeducation
- Normalize Symptoms
- Identify and manage triggers
- Draw connections
- Build Resilience
- Address Sleep Disturbances
- Build Trust
- Support empowerment
- Acknowledge Grief and Bereavement
- Use Culturally and Gender Responsive Services
- Make referrals where appropriate




# Trauma Specific Evidence-Based Practices

- Trauma Recovery Empowerment Model: TREM (male and female versions, as well as versions for adolescent girls and boys)
- Trauma Focused Cognitive Behavior Therapy (TF-CBT)
- Seeking Safety
- Mindfulness
- Eye Movement Desensitization and Reprocessing (EMDR)
- Exposure Therapy
- Sensory Interventions
- See EBP information on the SCCMHA website under Resources.

\*\*Most, if not all, of these interventions require specific training to practice.

EBP





## It Takes Everyone to Make a Difference

- Every contact with a consumer and with each other will affect us in 1 of 2 ways
  1. Contribute to a safe, trusting and healing environment
  2. Detract from a safe and trusting environment
- We all play a role in assisting consumers to heal and make progress in their lives
- We all matter when it comes to creating a safe, trusting, and healing environment
- How we treat an individual with a trauma history can shape how they view the service delivery system as a whole



SAGINAW COUNTY  
COMMUNITY MENTAL  
HEALTH AUTHORITY

# Preventing Re-traumatization



- Create a safe, welcoming environment, using consistency, communication, and compassion
- Support client-centered choices, ideals and autonomy—give as much personal choice and power as possible
- Always provide clear direct information about treatment and treatment stages



# What Helps

- Comfortable calming, and private treatment and waiting rooms
- Clean, comfortable furniture
- No “wrong door” philosophy
- Having a pleasant, hopeful demeanor when interacting with consumers
- A smile and a hello go a long way
- Be aware that trauma plays a role in everyone’s lives
- Interactions that express kindness, patience, reassurance, calm, acceptance and listening
- Clean, working, private restrooms
- Well-lit areas





# What Helps

- If there is security staff on the premises, they are visible yet approachable
- Recognize that an individual may be acting a certain way because they may be being triggered
- Sensible fair rules that focus on what you CAN do rather than what you CAN'T do
- Reacting in a calm way even if a consumer is acting out their emotions
- Viewing individuals as a whole person rather than their diagnosis or trauma
- Allowing someone to keep a door open or asking if you can close the door



# What Hurts


- Congested or noisy areas
- Unclean, uncomfortable areas
- Layers of red tape before a consumer's needs are met
- Questioning what is wrong with the person rather than what happened or how can I help
- Viewing an individual as their diagnosis rather than an individual who needs help.
- Ignoring that trauma exists
- Interactions that are humiliating, harsh, impersonal, disrespectful, critical, demanding or judgmental





SAGINAW COUNTY  
COMMUNITY MENTAL  
HEALTH AUTHORITY

# Moving from ACEs to Resilience



Moving from ACEs to RESILIENCE



## Important Resiliency Factors

- Realistic Optimism
- Facing fears
- Moral Compass
- Religion or spirituality
- Social Supports
- Resilient Role Models
- Physical fitness
- Brain fitness
- Cognitive and Emotional Flexibility
- Meaning and purpose
- Autonomy
- Self-esteem
- External supports
- Affiliation
- Safe adults



Burnout	Vicarious Trauma, Compassion Fatigue	Secondary Trauma, Indirect Trauma
Cumulative, usually over long period of time	Cumulative with symptoms that are unique to each service provider	Immediate and mirrors client/patient trauma
Predictable	Less predictable	Less predictable
Work dissatisfaction	Life dissatisfaction	Life dissatisfaction
Evident in work environment	Permeates work and home	Permeates work and home
Related to work environment conditions	Related to empathic relationship with <u>multiple</u> client's/patient's trauma experiences	Related to empathic relationship with one client's/patient's trauma experience
Can lead to health problems	Can lead to health problems	Can lead to health problems
Feel under pressure	Feel out of control	Feel out of control
Lack of motivation and/or energy	Symptoms of post-traumatic stress disorder	Symptoms of post-traumatic stress disorder similar to client/patient
No evidence of triggers	May have triggers that are unique to practitioner	Often have triggers that are similar to the client's/patient's triggers
Remedy is time away from work (vacation, stress leave) to recharge or positive change in work environment (this might mean a new job)	Remedy is treatment of self, similar to trauma treatment	Remedy is treatment of self, similar to trauma treatment



SAGINAW COUNTY  
COMMUNITY MENTAL  
HEALTH AUTHORITY

# Secondary Trauma and Self-care

- Hearing traumatic stories from consumers can cause secondary trauma in practitioners
- Recognize the signs of secondary trauma and get help when needed
- Supervisors should understand the signs and symptoms of secondary trauma
- Self-care practices should become a regular part of a practitioner's daily routine to help combat secondary trauma





## Signs and Symptoms of Secondary Trauma

- Hypervigilance
- Hopelessness
- Inability to embrace complexity
- Inability to listen and avoiding consumers
- Anger and cynicism
- Sleeplessness
- Fear
- Chronic exhaustion
- Physical ailments
- Minimizing
- Guilt



SAGINAW COUNTY  
COMMUNITY MENTAL  
HEALTH AUTHORITY

# What do you do for self-care?

"Almost everything will work  
again if you unplug it for a  
few minutes,  
including YOU."

-Anne Lamott

[www.paintedteacup.com](http://www.paintedteacup.com)



# Secondary Trauma Management/Intervention

- Psychoeducation
- Reflective clinical supervision
- Ongoing skills training
- Informal/self-report screening
- Workplace self-care groups or other form of peer-to-peer support and accountability
- Personal counseling
- Maintain balance with life and work
- Set clear boundaries with consumers
- Use of Evidence-Based Practices
- Exercise and good nutrition
- Mindfulness
- Grounding Exercises



# Helpful Resources

**Self-Help**  
These apps provide support and guidance in living with PTSD.

PTSD Coach   PTSD Family Coach   Beyond MST   Mindfulness Coach

**Treatment Companions**  
These apps offer additional help for PTSD treatments.

CPT Coach   PE Coach   CBT-i Coach   ACT Coach

**Related**  
These apps help with related issues affecting people with PTSD.

COVID Coach   Couples Coach   Insomnia Coach   StayQuit Coach


[Mobile Apps - PTSD: National Center for PTSD \(va.gov\)](https://www.va.gov/ncptsd/mobile-apps/)



SAGINAW COUNTY  
COMMUNITY MENTAL  
HEALTH AUTHORITY

# SCCMHA Resources

- [Trauma-Informed Care : Saginaw County Community Mental Health Authority \(sccmha.org\)](https://sccmha.org)
- [Trauma Screens : Saginaw County Community Mental Health Authority \(sccmha.org\)](https://sccmha.org)





# Grounding Exercise

Here's the 54321 “game”. This intervention can be used by anyone at anytime who is struggling with emotional dysregulation.

- Name 5 things you can see in the room with you.
- Name 4 things you can feel.
- Name 3 things you can hear right now.
- Name 2 things you can smell right now.
- Name 1 good thing about yourself.



SAGINAW COUNTY  
COMMUNITY MENTAL  
HEALTH AUTHORITY

# QUESTIONS?

[Mary.Baukus@sccmha.org](mailto:Mary.Baukus@sccmha.org)

989-272-7372



# References

- (2014). *Tip 57-Trauma Informed Care in Behavioral Health Situations* (First) [Review of *Tip 57-Trauma Informed Care in Behavioral Health Situations*]. HHS Publication. <https://store.samhsa.gov/sites/default/files/d7/priv/sma14-4816.pdf>
- American Psychiatric Association. (2023). *Stress & Trauma Toolkit for Treating LGBTQ in a Changing Political and Social Environment*. Psychiatry.org - LGBTQ. <https://www.psychiatry.org/psychiatrists/diversity/education/stress-and-trauma/lgbtq#:~:text=LGBTQ%20people%20who%20experience%20microtraumas,life%20and%20health%20care%20settings>.
- ClickPlayCEU.com (2015) Trauma-Informed Care. In *YouTube*. <https://www.youtube.com/watch?v=hJ4DbNWKmjo>
- Coping Skills*. (n.d.). Www.ibiblio.org. Retrieved November 5, 2019, from <http://www.ibiblio.org/rcip//copingskills.html>
- Dovetail Qld. (2019). Trauma and the Brain. In *YouTube*. [https://www.youtube.com/watch?v=ZLF\\_SEy6sdc](https://www.youtube.com/watch?v=ZLF_SEy6sdc)
- Floen, S. K., & Elklit, A. (2007). Psychiatric diagnoses, trauma, and suicidality. *Annals of General Psychiatry*. Retrieved October 13, 2022, from <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1858696/>
- Haro, B., Fernandez-Montalvo, J., Artega, A., & Lopez-Goni, J. J. (2023, September 15). *Gender Differences in Patients with Substance Use Disorder and Physical/Sexual Abuse: A Preliminary Study*. *Clínica y salud - journals.copmadrid.org*. [https://journals.copmadrid.org/clysa/archivos/1130\\_5274\\_clinsa\\_clysa2023a12.pdf](https://journals.copmadrid.org/clysa/archivos/1130_5274_clinsa_clysa2023a12.pdf)
- Home | Improving Mi Practices*. (n.d.). Www.improvingmipractices.org. Retrieved January 4, 2020, from <https://improvingmipractices.org>
- Kansas City PBS (2017). What is Trauma-Informed Care. In *YouTube*. <https://www.youtube.com/watch?v=AnBEROaeiak>



# References

- K. D. (2018, June 29). *What Is Complex Trauma?* The Awareness Centre. <https://theawarenesscentre.com/what-is-complex-trauma/>
- Michigan ACEs Data*. MDHHS - Prevalence of Trauma/Toxic Stress in Michigan. (n.d.). Retrieved January 4, 2020, from [https://www.michigan.gov/mdhhs/0,5885,7-339-73971\\_4911\\_69588\\_87676\\_87683---,00.html](https://www.michigan.gov/mdhhs/0,5885,7-339-73971_4911_69588_87676_87683---,00.html)
- Moving from ACEs to RESILIENCE*. (n.d.). Www.youtube.com. Retrieved December 16, 2019, from <https://www.youtube.com/watch?v=-pnhFmdz-ig>
- PTSD: National Center for PTSD (va.gov ). Mobile Apps, Retrieved April 26, 2022, <https://www.ptsd.va.gov/appvid/mobile/>
- Staff, J. S. (PACES C. (n.d.). *PACES Science 101 (FAQs) — Positive and Adverse Childhood Experiences*. PACESConnection. Retrieved November 5, 2019, from <https://www.acesconnection.com/blog/aces-101-faqs>
- The Jewish Federation. (2017). *What is trauma? - cdn.fedweb.org*. Center on Aging and Trauma. [https://cdn.fedweb.org/fed-42/2/AgingAndTrauma\\_FactSheet\\_CenterOnAgingAndTrauma\\_2fdbr.pdf](https://cdn.fedweb.org/fed-42/2/AgingAndTrauma_FactSheet_CenterOnAgingAndTrauma_2fdbr.pdf)
- The National Child Traumatic Stress Network*, (n.d.). The Learning Center, Retrieved December 16, 2021, from <https://www.nctsn.org/>.
- The National Council for Mental Well-Being*. (n.d.). Retrieved October 13, 2022, from <https://www.thenationalcouncil.org/>.
- The Trevor Project. (2022, July 28). *Trauma and suicide risk among LGBTQ youth*. NIHCM. <https://nihcm.org/publications/trauma-and-suicide-risk-among-lgbtq-youth>



SAGINAW COUNTY  
COMMUNITY MENTAL  
HEALTH AUTHORITY

Thank you



SAGINAW COUNTY  
COMMUNITY MENTAL  
HEALTH AUTHORITY

## **Main Facility**

500 Hancock, Saginaw, Michigan 48602

Phone: (989) 797-3400

Toll Free: 1-800-258-8678

Michigan Relay 711

## **24 Hour Mental Health Emergency Services**

(989) 792-9732

Toll Free: 1-800-233-0022

[www.sccmha.org](http://www.sccmha.org)

